


Infant mortality: includes deaths in the first year of age before the first birthday, including those died before the registration of birth

INFANT MORTALITY PER 1,000 LIVE BIRTHS



HOUSEHOLD EXPENDITURE FOR HEALTHCARE GOODS AND SERVICES IN 2013


Hospitalisations	211
Check-ups	124
Dentist	319
Auxiliary health services	164
Diagnostic tests	67
X-rays/ultrasound scans/Ecg	87
Glasses/contact lenses/blood pressure devices	178
Drugs	85
Thermometers/syringes	37

euro

6 HEALTH

DEATHS BY SOME CAUSES

1931, 1971 and 2011, per 100,000 inhabitants



NATIONAL HEALTH SERVICE INDICATORS


2011

	North	Centre	South and the Islands	Italy
General practitioners	19,608	9,771	16,682	46,061
General practitioner rate*	7.2	8.4	8.1	7.8
Patients per general practitioner	1,231	1,074	1,080	1,143
National Health Service paediatricians	3,258	1,517	2,941	7,716
National Health Service paediatrician rate**	8.6	9.6	9.6	9.2
Patients per paediatrician (under 14)	922	846	825	870
Non-emergency medical on-call services	795	416	1,670	2,881
On-call physicians	3,073	1,839	7,009	11,921
Local Health Authorities	74	29	42	145
Unified booking centres	65	29	41	135


* per 10,000 inhabitants ** per 10,000 children aged 0-14

HEALTH EXPENDITURE BY SECTOR

2010-2013, billions of euro at current prices



SMOKERS BY NUMBER OF DAILY CIGARETTES SMOKED IN 2013




Induced abortions: there has been a sharp reduction since Law n. 194/78 made induced abortions legal in Italy. 1982 and 1983 were the peak years, with a number of interventions exceeding 230,000 per year and an abortion rate of 16.7 and 16.4 respectively per 1,000 resident women aged 15 to 49. From that moment on, the number of induced abortions and the abortion rate progressively reduced to less than 8 per 1,000

Body Mass Index (BMI): determined by dividing the body weight in kilograms by the squared height in meters. A BMI under 18.5 is considered underweight, one of 18.5-24.99 is considered a healthy weight, one of 25-29.99 is overweight and one of 30 or more is obese

INDUCED ABORTIONS

1980-2011, per 1,000 resident women aged 15-49



DRUG CONSUMERS* BY AGE

1993-2013, thousands of people

	1993	1998	2007	2009	2011	2013
Up to 14	1,524	1,396	1,557	1,372	1,234	1,113
15-34	2,753	2,626	2,834	2,926	2,620	2,402
35-64	7,317	7,686	9,694	9,997	9,871	10,240
65 and over	5,986	6,933	9,008	9,509	9,607	10,318
Total	17,580	18,641	23,093	23,804	23,332	24,073

* in the two days preceding the interview

PERSONS BY BODY MASS INDEX

2013, per 100 persons aged 18 and over

